

1. Liu WM; **Chu WM**; Choudary PV; and Schmid CW (1995). Cell stress and translational inhibitors transiently increase the abundance of mammalian SINE transcripts. *Nuc. Acids Res.* 23, 1758-1765.
2. **Chu WM**; Liu WM; and Schmid CW (1995). RNA polymerase III promoter and terminator elements affect Alu RNA expression. *Nuc. Acids Res.* 23, 1750-1757.
3. Chesnokov I; **Chu WM**; and Schmid CW (1996). P53 inhibits RNA polymerase III-directed transcription in a promoter-dependent manner. *Mol. Cell. Biol.* 16, 7084-7088
4. **Chu WM**; Ballard R; and Schmid CW (1997). La protein and hairpin structure of termination elements facilitate Pol III transcription. *Nuc. Acids Res.* 25, 2077-2082.
5. **Chu WM**; Wang ZX; Roeder R.; and Schmid CW (1997). RNA polymerase III transcription repressed by Rb's interaction with TFIIB and TFIIC2. *J. Biol. Chem.* 272, 20072-78.
6. **Chu WM**; Ballard R; Carpick R; Williams BRG; and Schmid CW (1998). Potential Alu Function: Regulation of the activity of double-stranded RNA-activated kinase PKR. *Mol. Cell. Biol.* 18, 58-65.
7. Li ZW; **Chu WM**; Hu YL; Dalhase M; and Karin M. (1999) Role of IKK β in the activation of I κ B kinase and NF- κ B and cell survive defined by targeted gene disruption. *J Exp. Med.* 189(11), 1839-45.
8. **Chu WM**; Ostertag D; Li ZW; Chang.LF; Chen Y; Hu YL; Williams BRG, Perroult J and Karin M. (1999). JNK2 and IKK β are required for activating the innate response to viral infection. *Immunity* 11, 721.
9. **Chu WM**; Gong X; Li Z-W; Takabayashi K; Ouyang HH; Chen Y; Lois A; Chen DJ; Li GC; Karin M and Raz E (2000). DNA-PKcs is required for activation of innate immunity by immunostimulatory DNA. *Cell* 103, 901-9.
10. Dragoi AM; Fu X; Ivanov S; Zhang P; Shen L; Wu D; Li GC and **Chu WM** (2005). DNA-PKcs, but not TLR9 is required for activation of Akt by CpG-DNA. *EMBO J.* 24(4):779-89.
11. Zhang P, Dragoi AM, Chen J, Gong X, Ivanov S, Chuang T, Li Z-W, Tuthill C, Wan Y, Karin M and **Chu WM** (2005). Activation of IKK and NF- κ B by thymosin alpha 1 requires the TRAF6 and aPKC pathway. *EMBO Report*, 6(6): 531-7.
12. JP Xia, XZ Song, ZG Bi, **WM Chu**, YS Wan (2005). EGCG attenuates UV-induced apoptosis, NF κ B activation, and expression of proinflammatory cytokines and MMPs in cultured human keratinocytes *in vitro*. *Int J Mol Med.* 16(5):943-50.
13. XY Wang, ZG Bi, **WM Chu**, YS Wan (2005). IL-1 receptor antagonist inhibits MMP-1 production of UVA-irradiated human skin fibroblasts cultured with medium from UVB-irradiated keratinocytes. *Int J Mol Med.* 16(6):1117-24.